

## h-ITAC (AF647®), synthetic

**10 µg**

Catalogue number: CAF-13

### Almac Peptide and Protein Technologies

#### Chemokines

Custom Peptides

Site-Specific protein labelling

Modified Histones

Ubiquitylated peptides

### Product Information


<b>Product Name:</b>	h-ITAC (AF647®) CXCL11
<b>Synonyms:</b>	CXCL11(AF647®) human, Small-inducible cytokine B11; Interferon-inducible T-cell alpha chemoattractant; Interferon-gamma-inducible protein 9; IP-9; H174; Beta-R1
<b>Swiss Prot Accession No.:</b>	O14625
<b>Sequence:</b>	FPMFKRGRCL CIGPGVKAVK VADIEKASIM YPSNNCDKIE VIITLKENKG QRCLNPKSKQ ARLIIKKVER KNF-K(Alexa647®)-NH <sub>2</sub>
<b>Modifications:</b>	AF647® incorporated at indicated position in a molar ratio of dye:peptide of 1:1.

### Specification

<b>HPLC Purity:</b>	>95% area/area at 214nm
<b>Determined Mass:</b>	9.3 kDa
<b>Formulation / Appearance:</b>	Blue powder lyophilized from water/acetonitrile/ TFA to generate the TFA salt form. Peptide content determined by UV analysis at 280nm.

### Preparation and Storage

<b>Reconstitution / Storage:</b>	It is recommended that unopened vials are stored at -20 °C to -70 °C for periods of up to 18 months. Avoid repeat freeze-thaw cycles. It is recommended vials be centrifuged prior to opening. Water can be used to prepare stock solutions of 20 µmol.L <sup>-1</sup> . Stock solutions with up to 30% DMSO/water can also be prepared.
----------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------



*Not fully tested. For research use only. Not for use in human diagnostic or therapeutic procedures.*