

T +44 (0)28 3839 5717 • E almacshop@almacgroup.com • W shop.almacgroup.com

h-Fractalkine (AF647®), synthetic

10 µg

Catalogue number: CAF-14

Almac Peptide and Protein Technologies

Chemokines

Custom Peptides

Site-Specific protein labelling

Modified Histones

Ubiquitylated peptides

Product Information

Product h-Fractalkine (AF647®)

Name: CX3CL1

Synonyms: CX3CL1 (Alexa 647®) human, Neurotactin.

Swiss Prot P78423

Accession No:

Sequence: QHHGVTKCNI TCSKMTSKIP VALLIHYQQN QASCGKRAII

LETQRHRLFC ADPKEQWVKD AMQHLDRQAA ALTRNG-

Linker-K(Cys-AF647®)

Modifications: AF647® incorporated at indicated position in a molar ratio of

dye:peptide of 1:1. Linker used is 8-amino-3,6-dioxaoctanoic

acid and PEG thiazolidine.

Specification

HPLC Purity: >95% area/area at 214nm

Determined

Mass:

9.9 kDa

Formulation /

Appearance: the TI

the TFA salt form.

Peptide content determined by UV analysis at 280nm.

Preparation and Storage

Reconstitution / Storage:

It is recommended that unopened vials are stored at -20 °C to -70 °C for periods of up to 18 months. Avoid repeat freeze-thaw

Blue powder lyophilized from water/acetonitrile/TFA to generate

cycles.

It is recommended vials be centrifuged prior to opening.

Water can be used to prepare stock solutions of 20 μ mol.L-1. Stock solutions with up to 30% DMSO/water can also be

prepared.

Not fully tested. For research use only. Not for use in human diagnostic or therapeutic procedures.

