

SHOP

100 µg

h-TARC (Biotin), synthetic

Catalogue number: CB-16

Almac Peptide and Protein	Product Information	
Technologies Chemokines	Product Name:	h-TARC (Biotin) CCL17
Custom Peptides	Synonyms:	CCL17 human (Biotin), Small-inducible cytokine A17, Thymus and activation-regulated chemokine.
Site-Specific protein labelling	Curies Dest	, , , , , , , , , , , , , , , , , , ,
Modified Histones	Swiss Prot Accession No:	Q92583
Ubiquitylated peptides	Sequence:	ARGTNVGREC CLEYFKGAIP LRKLKTWYQT SEDCSRDAIV FVTVQGRAIC SDPNNKRVKN AVKYLQSLER SK(Linker- Biotin)
	Modifications:	Biotin incorporated at indicated position in a molar ratio of biotin:peptide of 1:1. Linker used is 8-amino-3,6-dioxaoctanoic acid.
	Specification	
	HPLC Purity:	>95% area/area at 214nm
	Determined Mass:	8.6 kDa
	Formulation / Appearance:	White powder lyophilized from water/acetonitrile/ TFA to generate the TFA salt form.
		Peptide content determined by UV analysis at 280nm.
	Preparation and	Storage
	Reconstitution / Storage:	It is recommended that unopened vials are stored at -20 °C to - 70 °C for periods of up to 18 months. Avoid repeat freeze-thaw cycles.
Sec Sec		It is recommended vials be centrifuged prior to opening.
Not fully te	ested. For research use o	Water can be used to prepare stock solutions of 20 µmol.L-1. Stock solutions with up to 30% DMSO/water can also be prepared. nly. Not for use in human diagnostic or therapeutic procedures.

Not fully tested. For research use only. Not for use in human diagnostic or therapeutic procedures.